
# **Evidence-Based Medicine**



# **Contents**

### **Purpose and procedure**

1

#### **EBM** notebook

- **2** Jottings ...
- 2 Delayed antibiotics for children with acute otitis media: is practice change sustainable?
- 4 Practice corner: should we screen all patients starting chemotherapy for chronic hepatitis B virus infection?
- **5** Evidence in context: one person's poison is another's acceptable risk
- **6** Letter: self-monitoring of blood glucose—caution warranted

# **Evidently**

7

# **Therapeutics**

- Review: medication adjustment based on fractional exhaled nitric oxide did not prevent asthma exacerbations
- 9 Benefits of early intensive glucose control to prevent diabetes complications were sustained for up to 10 years
- Benefits of tight blood pressure control to prevent diabetes complications were not sustained
- 11 A combination of telmisartan and ramipril increased renal outcomes in vascular disease or diabetes
- 12 General practitioners' advice to use topical rather than oral ibuprofen resulted in equivalent effects on chronic knee pain
- Alexander technique lessons were effective for chronic or recurrent back pain at 1 year
- 14 Review: daily salmeterol increases non-fatal serious adverse events in chronic asthma
- **15** Review: faecal occult blood test screening reduces risk of colorectal cancer mortality
- 16 CD4+ count—guided antiretroviral therapy interruption increased opportunistic disease and death more than continuous therapy in HIV
- 17 Active management of risk in pregnancy at term did not reduce caesarean delivery rates

# February 2009 Vol 14 No 1

- 18 Review: postmenopausal hormone therapy increases risk of stroke and venous thromboembolism but not coronary heart disease events
- 19 Review: an early invasive strategy has similar benefits in men and women with NSTEMI or unstable angina

# **Diagnosis**

20 A single measure of urinary neutrophil gelatinase—associated lipocalin was accurate for diagnosing acute kidney injury

# **Prognosis**

- 21 Review: people with epilepsy have higher risk of death by drowning than the general population
- Review: vasomotor symptoms peak at 1 year after final menstrual period and may persist for many years
- **23** Excess mortality after HIV seroconversion has greatly decreased in the past 10 years

### **Quality improvement**

Physician training in the use of a decision aid increased patient participation in decision making for CVD prevention

# **Clinical prediction guide**

- 25 The San Francisco Syncope Rule did not accurately predict serious short-term outcome in patients with syncope
- **26** The Paediatric Appendicitis Score (PAS) was useful in children with acute abdominal pain
- 27 The Splenoportal Index accurately predicted the presence of oesophageal varices in compensated cirrhosis

#### **Resource reviews**

28

# Additional articles abstracted in ACP Journal Club

30

#### Other articles noted

31

#### **Glossary**

Inside back cover

Evid Based Med: first published as on 1 February 2009. Downloaded from http://ebm.bmj.com/ on June 7, 2025 at Department GEZ-LTA Erasmushogeschool
Protected by copyright, including for uses related to text and data mining, Al training, and similar technologies.

BMJIJournals

This article has been chosen by the Editor to

This journal is a member of and subscribes to

the principles of the Committee on Publication

www.publicationethics.org.uk

be of special interest or importance and is

freely available online

COPE COMMITTEE ON PUBLICATION ETHICS