

Evidence-Based Medicine



Contents

Editorial

- 197** What evidence affects clinical practice? An analysis of Evidence-Based Medicine commentaries

EBM opinion and debate

- 198** Controversies in PSA screening

EBM Learning

- 199** Interpreting diagnostic tests with continuous results and no gold standard: a common scenario explained using the tuberculin skin test

- 202** Ten essential papers for the practice of evidence-based medicine

Research methods and reporting

- 205** What data sources do ophthalmologists trust?

Commentaries

Primary care

- 208** Routine prophylaxis is not necessary to prevent renal scarring in children with urinary tract infection

- 209** Supplementary antimicrobials for patients with HIV and <100 CD4 cells/ μ L are associated with improved survival

General medicine

- 210** Unblinded ASCOT study results do not rule out that muscle symptoms are an adverse effect of statins

- 211** SOFA criteria predict infection-related in-hospital mortality in ICU patients better than SIRS criteria and the qSOFA score

- 212** Prescribing antibiotics to hospitalised patients increases the risk of *Clostridium difficile* infection for the next bed occupant

- 213** Cephalixin plus trimethoprim-sulfamethoxazole was not superior to cephalixin alone for the treatment of outpatient non-purulent cellulitis

- 214** Adjunctive antibiotics for drained skin abscesses improve clinical cure rate

- 215** Ocrelizumab appears to reduce relapse and disability in multiple sclerosis but quality of evidence is moderate

- 217** In patients with localised prostate cancer, active surveillance is associated with better sexual function, urinary symptoms and bowel symptoms

December 2017 Volume 22 Issue 6

- 218** A meta-analysis of positive airway pressure treatment for cardiovascular prevention: why mix apples and pears?

- 219** Intensive glucose control in patients with diabetes prevents onset and progression of microalbuminuria, but effects on end-stage kidney disease are still uncertain

Surgery

- 221** Don't put off until tomorrow what you can do today: Early cholecystectomy is cost-effective in symptomatic cholelithiasis requiring hospitalization

- 222** Minimal difference in survival between radical prostatectomy and observation in men with modest life expectancy

Emergency care

- 223** Protocolised early goal-directed therapy in patients with sepsis/septic shock does not result in improved survival compared with usual care with less invasive resuscitation strategies

- 224** Stent choice in cardiogenic shock complicating acute myocardial infarction likely does not affect mortality or reinfarction

- 225** Prophylactic platelet transfusion does not reduce risk of clinical bleeding in adults with dengue and thrombocytopaenia

- 226** Acute myocardial infarction can be ruled out with a single high-sensitivity cardiac troponin T level

- 227** Early invasive strategy should be performed within 72 hours in high-risk patients with non-ST-elevation myocardial infarction

Maternal and child health

- 228** Long-acting reversible contraception acceptability and satisfaction is high among adolescents


- 229** Acupuncture is not as effective as infertility treatment in women with PCOS


Journal club

- 231** Journal Clubs: 1. Origins

- 232** Journal Clubs: 2. Why and how to run them and how to publish them

- 235** Evaluation of a 0-hour/1-hour algorithm in the diagnosis of myocardial infarction with high-sensitivity cardiac troponin T1

 This article has been chosen by the Editor to be of special interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebm.bmj.com/site/about/guidelines.xhtml#open>

 This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it