


# Evidence-Based Medicine

**Editors:** Brian Haynes & Paul Glasziou



**Contents**

Purpose and procedure	130
Additional articles abstracted in <i>ACP Journal Club</i>	131

**EBM notebook**

Practice corner: using clinical prediction rules	132
Resource review	135

**Therapeutics**

Oestrogen plus progestogen increased coronary heart disease and breast cancer events in postmenopausal women	137
Review: postmenopausal oestrogen replacement therapy increases the risk for venous thromboembolism	138
A lifestyle intervention or metformin prevented or delayed the onset of type 2 diabetes in people at risk	139
Warfarin combined with low dose aspirin in myocardial infarction did not provide clinical benefit beyond that of aspirin alone	140
Review: angiotensin receptor blockers do not reduce mortality or hospital admission rates in heart failure	141
Review: n-3 polyunsaturated fatty acids reduce fatal myocardial infarction, sudden death, and overall mortality in coronary artery disease	142
Parental education and guided self management of asthma or wheeze in pre-school children was not effective	143
Review: inhaled corticosteroids for asthma or chronic obstructive pulmonary disease do not increase bone loss	144
Review: nedocromil sodium decreases airflow reduction and recovery time in exercise induced bronchoconstriction	145
Review: $\beta$ 2 agonists are ineffective but increase adverse effects in acute bronchitis without underlying pulmonary disease	146
Review: tricyclic antidepressants, capsaicin, gabapentin, and oxycodone are effective for postherpetic neuralgia	147

Review: home visiting with multidimensional assessment and multiple visits is effective in low risk elderly people	148
Sertraline maintenance treatment reduced relapse and dropouts in post-traumatic stress disorder	149
Review: brief interventions reduce drinking in patients not seeking treatment	150
A brief intervention reduced alcohol drinking for up to 48 months in problem drinkers	151
A normal calcium, low protein, low salt diet reduced recurrence of renal stones over 5 years	152
Physiotherapy or a wait and see policy were the best options for lateral epicondylitis at 1 year	153
Bed rest, physiotherapy, and continuation of activities of daily living were equally effective for sciatica	154
Review: antiemetic drugs reduce nausea in early pregnancy	155
Telephone-based peer support increased the duration of breast feeding in primiparous mothers	156
Clonidine and methylphenidate were effective for attention deficit hyperactivity disorder in children with comorbid tics	157
Pharmacist review changed more repeat prescriptions for elderly patients than usual general practice review	158

**Diagnosis**

Review: questionnaires for detecting clinical depression in primary care have similar diagnostic accuracy	159
---	-----

**Aetiology**

Increased plasma total homocysteine levels increased the risk for dementia in the elderly	160
---	-----